



Splint Safety & Care

Splint Cleaning: We recommend brushing your retainers under cool water with a toothbrush and toothpaste when you brush your teeth. For disinfecting, soak the retainer 2X/week for no more than 15 minutes. Your retainer can be soaked in any denture or retainer cleaner and Listerine.

Safety: If the retainers are lost or damaged beyond repair, we will charge a fee for replacement. Also, call us as soon as possible to prevent your teeth from shifting.

To avoid this, please follow these rules:

- Do not leave your retainer on a cafeteria tray, or wrap it in a napkin or paper towel; you may accidentally discard it. Use the proper case provided for storage of your retainer when it is not in your mouth.
- Do not leave your retainer in a place where animals can get them as they enjoy chewing on them.
- Do not put your retainer in your pocket as this may cause the wires to become bent or broken.
- Do not place the retainer in hot water or near heaters.
- Do not chew on your retainer. This will flatten them and they will no longer fit.

Adjusting to your splint: You may find it difficult to speak clearly at first, however, with practice your speech will return to normal. Also, for several days after the retainers are placed, extra saliva may be produced. This will gradually lessen.

We invite you to call us if you have any concerns or if you would just like to say HELLO!

Thank you, and best wishes for a wonderful life!

Dr. Gary Cartwright and Staff